

Zn-SURE is a
Zinc solution
chelated with
Fulvic Acids



BENEFITS

Registered under Act 36 of 1947

Nutrition: Zinc plays a vital role in chlorophyll and carbohydrate production.

Zinc is responsible for the pH regulation in the cytoplasm of cells.

Zn-Sure increases the availability of the zinc nutrient, preventing the mineral from forming insoluble precipitates, reducing toxicity of some metal ions in the plant, preventing the nutrient from leaching and increasing the mobility of the nutrient in the plant.

Zinc assists in the regulation of phosphorus uptake in the plant.

PACKAGING

5 L, 25 L, 1000 L
Product Code: ZN01





APPLICATION RATES

FOLIAR SPRAY:

Vegetables & Leafy Greens

Apply as a 1% spray solution (dissolve 1 L Zn-Sure in 100 Litres of water) at an application rate of up to 5 Litres Zn-Sure per hectare (Up to 500 litres/hectare of a 1% solution) as a nutrient source. Apply at bud before flowering, then repeat one month after set. For Leafy greens: apply at 5th leaf, repeat in 2 weeks.

Orchards & Vineyards

Application rates as above. Apply between bud and flowering, after fruit set and once during fruit-fill or as required.

Ornamentals & Turf

Application rates as above. Apply every 2 weeks or as required.

Broad acre Crops

300 mL – 1 L/ha Zn-Sure in 60 – 100 L water, depending on canopy closure. Do not exceed a concentration of 1 L Zn-Sure per 100 L water. Apply once or twice prior to flowering.

Pasture

300 mL – 1 L/ha Zn-Sure in 60 – 100 L water. Do not exceed a concentration of 1 L Zn-Sure per 100 L water. Apply as required.

Spot-Spraying

Dilute 50 mL Zn-Sure per 10 L water. Apply 200ml to 400ml solution per square meter.

FERTIGATION:

5 - 10 L/ha as a nutrient source.

HOME GARDEN APPLICATION:

Dilute 15 mL Zn-Sure in a 9 L watering can and apply the diluted mix at a rate of 1 litre per square metre. Apply fortnightly or as required

COMPANY DETAILS:

Wolseley Liquids CC

(reg. no. 2010/018108/23)

57 Euufees street, Wolseley, 6830. Western Cape Province, South Africa

Phone +27 23 231 0454

Fax +27 86 768 0374

admin@wolseleyliquids.com

www.wolseleyliquids.com

TYPICAL ANALYSIS W/V

Total Zinc	67.2g/kg
SG	1.15
pH	1.12
Appearance	Translucent Liquid

INSTRUCTIONS

Shake or stir well before use. Where higher water rates are required for good foliar coverage, do not exceed maximum product rate per hectare. Sensitive foliage or plants should be test sprayed prior to full scale application. Frequencies are a guideline only and can be adjusted according to farm practices and/or in conjunction with crop monitoring. Ensure suitable spray conditions (e.g. temperature, humidity, wind speed), prior to foliar application. Compatibility and/or performance cannot be guaranteed when combined with other products. Jar test for compatibility and spray test on small area before wide-spread use.

STORAGE & HANDLING

Store in a cool dry area out of direct sunlight. This product has been tested to store safely down to 5°C. Crystallisation or sedimentation may occur below 5°C. Do not pre-mix or store in diluted form. Seal lid/cap immediately after use to prevent contamination of airborne microbes.

Disclaimer: Any recommendations provided by 'Wolseley Liquids CC' or its Distributors are advice only. As no control can be exercised over storage, handling, mixing, application or use, or weather, plant or soil conditions before, during or after application (all of which may affect the performance of our program), no responsibility for, or liability for any failure in performance, losses, damages, or injuries (consequential or otherwise), arising from such storage, mixing, application, or use will be accepted under any circumstances whatsoever. The Buyer assumes all responsibility for the use of any 'Wolseley Liquids CC' products.

DISTRIBUTOR INFO: